arXiv:hep-ph/0504223v2 29 Nov 2005

A new method for calculation of traces of Dirac γ-matrices in Minkowski space

Alexander L. Bondarev
National Scientific and Educational Center of Particle and High Energy Physics of the Belarusian State University
e-mail: bondarev@hep.by

Abstract

This paper presents some relations for orthonormal bases in the Minkowski space and isotropic tetrads constructed from the vectors of these bases.

As an example of an application of the obtained formulae, in particular recursion relations, a new method is proposed to calculate traces of Dirac γ-matrices in the Minkowski space. Compared to the classical algorithms, the new method results in more compact expressions for the traces. Specifically, it may be easily implemented as a simple yet efficient computer algorithm.

1 Introduction

The problems arising when the traces of a large number of Dirac γ-matrices should be calculated are well known. For ten matrices and more the expressions for the traces become extremely cumbersome.

For example, the classical algorithm based on the application of the algebra of γ-matrices

$$\gamma^{\mu}\gamma^{\nu} + \gamma^{\nu}\gamma^{\mu} = 2g^{\mu\nu} \tag{1}$$

for the trace of ten γ-matrices results in the expression consisting of 945 terms (e.g. such algorithm is used in REDUCE [1]).

In four-dimensional space-time somewhat shorter results provides the method using the Chisholm identities [2] and Kahane algorithm [3] that is based on the reduction formula

$$\gamma^{\mu}\gamma^{\nu}\gamma^{\rho} = g^{\mu\nu}\gamma^{\rho} - g^{\mu\rho}\gamma^{\nu} + g^{\nu\rho}\gamma^{\mu} + i\varepsilon^{\mu\nu\rho\lambda}\gamma^{5}\gamma_{\lambda}\ . \tag{2}$$

This method, implemented first in SCHOONSCHIP [4] and later in FORM [5], gives for the trace of ten γ-matrices the following expression consisting of 693 terms:

$$\mathrm{Tr}_4(\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6\hat{a}_7\hat{a}_8\hat{a}_9\hat{a}_{10}) = \mathrm{Tr}(\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6\hat{a}_7\hat{a}_8\hat{a}_9\hat{a}_{10})$$

$$+\ G\begin{pmatrix} a_1 & a_2 & a_3 & a_6 & a_7 \\ a_4 & a_5 & a_8 & a_9 & a_{10} \end{pmatrix} \ -\ G\begin{pmatrix} a_1 & a_2 & a_3 & a_8 & a_9 \\ a_4 & a_5 & a_6 & a_7 & a_{10} \end{pmatrix}$$
$$+\ G\begin{pmatrix} a_1 & a_2 & a_3 & a_8 & a_{10} \\ a_4 & a_5 & a_6 & a_7 & a_9 \end{pmatrix} \ -\ G\begin{pmatrix} a_1 & a_2 & a_3 & a_9 & a_{10} \\ a_4 & a_5 & a_6 & a_7 & a_8 \end{pmatrix} \quad .$$

The proposed new method results in more compact expressions (e.g. consisting of 32 terms for the trace of ten γ-matrices) than the classical methods. In particular, this method may be easily implemented as a simple yet efficient computer algorithm.

The method is based on the properties of orthonormal bases in the Minkowski space and isotropic tetrads (see e.g. [6]) constructed from the vectors of these bases.

In this paper we use the Feynman metrics:

$$\mu = 0,1,2,3, \quad a^\mu = (a_0, \vec{a}), \quad a_\mu = (a_0, -\vec{a}), \quad (ab) = a_\mu b^\mu = a_0 b_0 - \vec{a}\vec{b}\ ,$$

and a sign of the Levi-Civita tensor is determined as $\varepsilon_{0123} = +1$;

$$\hat{a} = \gamma_\mu a^\mu\ , \ \ \gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3\ .$$

Together with the regular Gram determinants, we use the generalized ones. For example:

$$G\begin{pmatrix} \mu & \alpha \\ \nu & \beta \end{pmatrix} = \begin{vmatrix} g^{\mu\nu} & g^{\mu\beta} \\ g^{\alpha\nu} & g^{\alpha\beta} \end{vmatrix} = g^{\mu\nu}g^{\alpha\beta} - g^{\mu\beta}g^{\alpha\nu}, \ \ g^{\mu\nu} = \begin{cases} \ \ 1, & \mu = \nu = 0 \\ -1, & \mu = \nu = 1,2,3 \\ \ \ 0, & \mu \neq \nu \end{cases}$$

$$G\begin{pmatrix} \mu & \alpha \\ \nu & \beta \end{pmatrix} a_\alpha b_\beta = G\begin{pmatrix} \mu & a \\ \nu & b \end{pmatrix} = \begin{vmatrix} g^{\mu\nu} & b^\mu \\ a^\nu & (ab) \end{vmatrix} = g^{\mu\nu}(ab) - b^\mu a^\nu\ \ .$$

2 Some relations for isotropic tetrads in the Minkowski space

Let vectors l_0, l_1, l_2, l_3 form an orthonormal basis in the Minkowski space, i.e.

$$(l_A l_B) = g_{AB}\ , \ \ (A, B = 0,1,2,3)$$

(index of the basis vector is denoted by the capital letters).

In four-dimensional space both the regular and generalized fifth-order Gram determinants equal zero, therefore

$$G\begin{pmatrix} \mu & l_0 & l_1 & l_2 & l_3 \\ \nu & l_0 & l_1 & l_2 & l_3 \end{pmatrix} = \begin{vmatrix} g^{\mu\nu} & l_0^\mu & l_1^\mu & l_2^\mu & l_3^\mu \\ l_0^\nu & +1 & 0 & 0 & 0 \\ l_1^\nu & 0 & -1 & 0 & 0 \\ l_2^\nu & 0 & 0 & -1 & 0 \\ l_3^\nu & 0 & 0 & 0 & -1 \end{vmatrix}$$
$$= -g^{\mu\nu} + l_0^\mu l_0^\nu - l_1^\mu l_1^\nu - l_2^\mu l_2^\nu - l_3^\mu l_3^\nu = 0\ . \qquad (3)$$

Whence we obtain the expression for the metric tensor:

$$g^{\mu\nu} = l_0^\mu l_0^\nu - l_1^\mu l_1^\nu - l_2^\mu l_2^\nu - l_3^\mu l_3^\nu \ . \tag{4}$$

Furthermore, for the orthonormal basis we have

$$G\begin{pmatrix} l_0 & l_1 & l_2 & l_3 \\ l_0 & l_1 & l_2 & l_3 \end{pmatrix} = \begin{vmatrix} +1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{vmatrix} = -1 \ . \tag{5}$$

Since

$$G\begin{pmatrix} l_0 & l_1 & l_2 & l_3 \\ l_0 & l_1 & l_2 & l_3 \end{pmatrix} = -\varepsilon_{\mu\nu\rho\lambda} l_0^\mu l_1^\nu l_2^\rho l_3^\lambda \ \varepsilon_{\alpha\beta\sigma\tau} l_0^\alpha l_1^\beta l_2^\sigma l_3^\tau \ , \tag{6}$$

we get:

$$\varepsilon_{\mu\nu\rho\lambda} l_0^\mu l_1^\nu l_2^\rho l_3^\lambda = \pm 1 \ . \tag{7}$$

The "+" and "−" signs correspond to the "right-hand" and "left-hand" quadruples of the vectors.

For definiteness in the following we consider the "right-hand" basis.

$$\begin{aligned} \hat{l}_0\hat{l}_1\hat{l}_2\hat{l}_3 &= \frac{1}{4}\left[\mathrm{Tr}\left(\hat{l}_0\hat{l}_1\hat{l}_2\hat{l}_3\right) + \gamma^5\,\mathrm{Tr}\left(\gamma^5\hat{l}_0\hat{l}_1\hat{l}_2\hat{l}_3\right) + \frac{1}{2}\sigma^{\mu\nu}\,\mathrm{Tr}\left(\sigma_{\mu\nu}\hat{l}_0\hat{l}_1\hat{l}_2\hat{l}_3\right)\right] \\ &= \frac{1}{4}\gamma^5\,\mathrm{Tr}\left(\gamma^5\hat{l}_0\hat{l}_1\hat{l}_2\hat{l}_3\right) = i\gamma^5\varepsilon_{\mu\nu\rho\lambda} l_0^\mu l_1^\nu l_2^\rho l_3^\lambda = i\gamma^5 \ , \end{aligned} \tag{8}$$

which gives us additional relations for the basis vectors, for example:

$$\hat{l}_0 = i\gamma^5\hat{l}_1\hat{l}_2\hat{l}_3 \ , \ \ \hat{l}_1 = i\gamma^5\hat{l}_0\hat{l}_2\hat{l}_3 \ , \ \ \hat{l}_2 = -i\gamma^5\hat{l}_0\hat{l}_1\hat{l}_3 \ , \ \ \hat{l}_3 = i\gamma^5\hat{l}_0\hat{l}_1\hat{l}_2 \ . \tag{9}$$

Using (9), we derive the expressions which we need in the following:

$$\hat{l}_0\hat{l}_1 = -i\gamma^5\hat{l}_2\hat{l}_3 \ , \tag{10}$$

$$\hat{l}_0\hat{l}_2 = i\gamma^5\hat{l}_1\hat{l}_3 \ , \tag{11}$$

$$\hat{l}_0\hat{l}_3 = -i\gamma^5\hat{l}_1\hat{l}_2 \ , \tag{12}$$

$$\hat{l}_1\hat{l}_2 = i\gamma^5\hat{l}_0\hat{l}_3 \ , \tag{13}$$

$$\hat{l}_1\hat{l}_3 = -i\gamma^5\hat{l}_0\hat{l}_2 \ , \tag{14}$$

$$\hat{l}_2\hat{l}_3 = i\gamma^5\hat{l}_0\hat{l}_1 \ . \tag{15}$$

Let us introduce the isotropic tetrad

$$q_+ = \frac{l_0 + l_1}{\sqrt{2}} \ , \ \ q_- = \frac{l_0 - l_1}{\sqrt{2}} \ , \ \ (q_+ q_-) = 1 \ ; \tag{16}$$

$$e_+ = \frac{l_2 + il_3}{\sqrt{2}} \ , \quad e_- = \frac{l_2 - il_3}{\sqrt{2}} \ , \quad (e_+ e_-) = -1 \ , \quad (e_\mp)^* = e_\pm \ ; \tag{17}$$

$$(q_+ e_+) = (q_+ e_-) = (q_- e_+) = (q_- e_-) = 0 \ .$$

Then (4) takes the form

$$g^{\mu\nu} = q_+^\mu q_-^\nu + q_-^\mu q_+^\nu - e_+^\mu e_-^\nu - e_-^\mu e_+^\nu \ , \tag{18}$$

from which follow the expressions below:

$$(ab) = (aq_+)(bq_-) + (aq_-)(bq_+) - (ae_+)(be_-) - (ae_-)(be_+) \ , \tag{19}$$

$$\gamma^\mu = q_+^\mu \hat{q}_- + q_-^\mu \hat{q}_+ - e_+^\mu \hat{e}_- - e_-^\mu \hat{e}_+ \ , \tag{20}$$

$$\hat{a} = (aq_+)\hat{q}_- + (aq_-)\hat{q}_+ - (ae_+)\hat{e}_- - (ae_-)\hat{e}_+ \ . \tag{21}$$

From (10) – (15) we have

$$\begin{aligned} \hat{q}_+ \hat{e}_- &= \frac{1}{2}(\hat{l}_0 + \hat{l}_1)(\hat{l}_2 - i\hat{l}_3) = \frac{1}{2}(\hat{l}_0\hat{l}_2 + \hat{l}_1\hat{l}_2 - i\hat{l}_0\hat{l}_3 - i\hat{l}_1\hat{l}_3) \\ &= \frac{1}{2}(i\gamma^5\hat{l}_1\hat{l}_3 + i\gamma^5\hat{l}_0\hat{l}_3 - \gamma^5\hat{l}_1\hat{l}_2 - \gamma^5\hat{l}_0\hat{l}_2) \\ &= -\gamma^5\frac{1}{2}(\hat{l}_0\hat{l}_2 + \hat{l}_1\hat{l}_2 - i\hat{l}_0\hat{l}_3 - i\hat{l}_1\hat{l}_3) = -\gamma^5\hat{q}_+\hat{e}_- \ . \end{aligned} \tag{22}$$

In a similar manner

$$\hat{q}_-\hat{e}_+ = -\gamma^5\hat{q}_-\hat{e}_+ \, . \tag{23}$$

$$\hat{q}_+\hat{e}_+ = \gamma^5\hat{q}_+\hat{e}_+ \, . \tag{24}$$

$$\hat{q}_-\hat{e}_- = \gamma^5\hat{q}_-\hat{e}_- \, . \tag{25}$$

From (22) – (25) it follows that

$$(1+\gamma^5)\hat{q}_+\hat{e}_- = 0 \ , \tag{26}$$

$$(1+\gamma^5)\hat{q}_-\hat{e}_+ = 0 \ , \tag{27}$$

$$(1-\gamma^5)\hat{q}_+\hat{e}_+ = 0 \ , \tag{28}$$

$$(1-\gamma^5)\hat{q}_-\hat{e}_- = 0 \ . \tag{29}$$

Hence

$$(1+\gamma^5)\hat{e}_-\hat{e}_+\hat{q}_- = -2(1+\gamma^5)\hat{q}_- \ , \tag{30}$$

$$(1+\gamma^5)\hat{e}_+\hat{e}_-\hat{q}_+ = -2(1+\gamma^5)\hat{q}_+ \ , \tag{31}$$

$$(1+\gamma^5)\hat{q}_-\hat{q}_+\hat{e}_- = 2(1+\gamma^5)\hat{e}_- \ , \tag{32}$$

$$(1+\gamma^5)\hat{q}_+\hat{q}_-\hat{e}_+ = 2(1+\gamma^5)\hat{e}_+ \ . \tag{33}$$

$$(1-\gamma^5)\hat{e}_+\hat{e}_-\hat{q}_- = -2(1-\gamma^5)\hat{q}_- \ , \tag{34}$$

$$(1-\gamma^5)\hat{e}_-\hat{e}_+\hat{q}_+ = -2(1-\gamma^5)\hat{q}_+ \ , \tag{35}$$

$$(1-\gamma^5)\hat{q}_+\hat{q}_-\hat{e}_- = 2(1-\gamma^5)\hat{e}_- \ , \tag{36}$$

$$(1-\gamma^5)\hat{q}_-\hat{q}_+\hat{e}_+ = 2(1-\gamma^5)\hat{e}_+ \ . \tag{37}$$

For example, for (30) we have

$$(1+\gamma^5)\hat{e}_-\hat{e}_+\hat{q}_- = 2(e_-e_+)(1+\gamma^5)\hat{q}_- - (1+\gamma^5)\hat{e}_+\hat{e}_-\hat{q}_- \ ,$$

where the second term on the right-hand side equals zero due to (29).

Furthermore, from the definition of the isotropic tetrad (16), (17) it is clear that

$$\hat{q}_-\hat{q}_+\hat{q}_- = 2\hat{q}_- \ , \tag{38}$$

$$\hat{q}_+\hat{q}_-\hat{q}_+ = 2\hat{q}_+ \ , \tag{39}$$

$$\hat{e}_-\hat{e}_+\hat{e}_- = -2\hat{e}_- \ , \tag{40}$$

$$\hat{e}_+\hat{e}_-\hat{e}_+ = -2\hat{e}_+ \ . \tag{41}$$

It is important, that any three operators of the form $\hat{q}_\lambda$ or $\hat{e}_\sigma$ taken in succession in the presence of $(1 \pm \gamma^5)$ are either reduced to only one of them or equal zero.

3 Recursion relations

Now, using (21), (26) – (41) we have

$$\begin{aligned}
&(1+\gamma^5)\hat{q}_-\hat{a}\hat{b} = (aq_-)(1+\gamma^5)\hat{q}_-\hat{q}_+\hat{b} - (ae_+)(1+\gamma^5)\hat{q}_-\hat{e}_-\hat{b} \\
= \ & (aq_-)[(bq_+)(1+\gamma^5)\hat{q}_-\hat{q}_+\hat{q}_- - (be_+)(1+\gamma^5)\hat{q}_-\hat{q}_+\hat{e}_-] \\
- \ & (ae_+)[(bq_-)(1+\gamma^5)\hat{q}_-\hat{e}_-\hat{q}_+ - (be_-)(1+\gamma^5)\hat{q}_-\hat{e}_-\hat{e}_+] \\
= \ & (aq_-)2[(bq_+)(1+\gamma^5)\hat{q}_- - (be_+)(1+\gamma^5)\hat{e}_-] \\
- \ & (ae_+)2[-(bq_-)(1+\gamma^5)\hat{e}_- + (be_-)(1+\gamma^5)\hat{q}_-] \\
= \ & 2[(aq_-)(bq_+) - (ae_+)(be_-)](1+\gamma^5)\hat{q}_- \\
+ \ & 2[(ae_+)(bq_-) - (aq_-)(be_+)](1+\gamma^5)\hat{e}_- \ .
\end{aligned} \tag{42}$$

In the same way

$$\begin{aligned}
&(1+\gamma^5)\hat{e}_-\hat{a}\hat{b} = (aq_+)(1+\gamma^5)\hat{e}_-\hat{q}_-\hat{b} - (ae_-)(1+\gamma^5)\hat{e}_-\hat{e}_+\hat{b} \\
= \ & (aq_+)[(bq_-)(1+\gamma^5)\hat{e}_-\hat{q}_-\hat{q}_+ - (be_-)(1+\gamma^5)\hat{e}_-\hat{q}_-\hat{e}_+]
\end{aligned}$$

$$
\begin{aligned}
- \quad & (ae_-)[(bq_+)(1+\gamma^5)\hat{e}_-\hat{e}_+\hat{q}_- - (be_+)(1+\gamma^5)\hat{e}_-\hat{e}_+\hat{e}_-] \\
= \quad & (aq_+)2[(bq_-)(1+\gamma^5)\hat{e}_- - (be_-)(1+\gamma^5)\hat{q}_-] \\
- \quad & (ae_-)2[-(bq_+)(1+\gamma^5)\hat{q}_- + (be_+)(1+\gamma^5)\hat{e}_-] \\
= \quad & 2[(aq_+)(bq_-) - (ae_-)(be_+)](1+\gamma^5)\hat{e}_- \\
+ \quad & 2[(ae_-)(bq_+) - (aq_+)(be_-)](1+\gamma^5)\hat{q}_- \; . \qquad (43)
\end{aligned}
$$

Defining the functions

$$F_1(a,b) = 2[(aq_-)(bq_+) - (ae_+)(be_-)] \; , \qquad (44)$$

$$F_2(a,b) = 2[(ae_+)(bq_-) - (aq_-)(be_+)] = 2 \; G\begin{pmatrix} a & b \\ e_+ & q_- \end{pmatrix} \; , \qquad (45)$$

$$F_3(a,b) = 2[(aq_+)(bq_-) - (ae_-)(be_+)] \; , \qquad (46)$$

$$F_4(a,b) = 2[(ae_-)(bq_+) - (aq_+)(be_-)] = 2 \; G\begin{pmatrix} a & b \\ e_- & q_+ \end{pmatrix} \; , \qquad (47)$$

we finally get

$$(1+\gamma^5)\hat{q}_-\hat{a}\hat{b} = F_1(a,b)(1+\gamma^5)\hat{q}_- + F_2(a,b)(1+\gamma^5)\hat{e}_- \; , \qquad (48)$$

$$(1+\gamma^5)\hat{e}_-\hat{a}\hat{b} = F_3(a,b)(1+\gamma^5)\hat{e}_- + F_4(a,b)(1+\gamma^5)\hat{q}_- \; . \qquad (49)$$

Making similar calculations, we obtain

$$(1+\gamma^5)\hat{q}_+\hat{a}\hat{b} = F_3(a,b)(1+\gamma^5)\hat{q}_+ + F_4(a,b)(1+\gamma^5)\hat{e}_+ \; , \qquad (50)$$

$$(1+\gamma^5)\hat{e}_+\hat{a}\hat{b} = F_1(a,b)(1+\gamma^5)\hat{e}_+ + F_2(a,b)(1+\gamma^5)\hat{q}_+ \; . \qquad (51)$$

In a similar manner, after defining the functions

$$F_5(a,b) = 2[(aq_-)(bq_+) - (ae_-)(be_+)] \; , \qquad (52)$$

$$F_6(a,b) = 2[(ae_-)(bq_-) - (aq_-)(be_-)] = 2 \; G\begin{pmatrix} a & b \\ e_- & q_- \end{pmatrix} \; , \qquad (53)$$

$$F_7(a,b) = 2[(aq_+)(bq_-) - (ae_+)(be_-)] \; , \qquad (54)$$

$$F_8(a,b) = 2[(ae_+)(bq_+) - (aq_+)(be_+)] = 2 \; G\begin{pmatrix} a & b \\ e_+ & q_+ \end{pmatrix} \qquad (55)$$

we obtain

$$(1-\gamma^5)\hat{q}_-\hat{a}\hat{b} = F_5(a,b)(1-\gamma^5)\hat{q}_- + F_6(a,b)(1-\gamma^5)\hat{e}_+ \; , \qquad (56)$$

$$(1-\gamma^5)\hat{e}_+\hat{a}\hat{b} = F_7(a,b)(1-\gamma^5)\hat{e}_+ + F_8(a,b)(1-\gamma^5)\hat{q}_- \; , \qquad (57)$$

$$(1-\gamma^5)\hat{q}_+\hat{a}\hat{b} = F_7(a,b)(1-\gamma^5)\hat{q}_+ + F_8(a,b)(1-\gamma^5)\hat{e}_- \ , \tag{58}$$

$$(1-\gamma^5)\hat{e}_-\hat{a}\hat{b} = F_5(a,b)(1-\gamma^5)\hat{e}_- + F_6(a,b)(1-\gamma^5)\hat{q}_+ \ . \tag{59}$$

Note that the functions defined in (44) – (47), (52) – (55) are not independent:

$$F_5(a,b) = F_1^*(a,b) \ ,$$

$$F_6(a,b) = F_2^*(a,b) \ ,$$

$$F_7(a,b) = F_3^*(a,b) \ ,$$

$$F_8(a,b) = F_4^*(a,b) \ .$$

Furthermore, the following relations hold:

$$F_3(b,a) = F_1(a,b) \ ,$$

$$F_1(a,b) + F_3(a,b) = 2(ab) \ ,$$

$$F_1(a,b)F_3(a,b) - F_2(a,b)F_4(a,b) = a^2b^2$$

(see also **Appendix B**).

4 A new method for calculation of traces

We now use the formulae (44) – (59) obtained in the previous section to calculate the traces of Dirac γ-matrices as follows:

$$\begin{aligned}\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_1\hat{a}_2\hat{a}_3\cdots\hat{a}_{2n+1}] &= F_1(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_3\cdots\hat{a}_{2n+1}] \\ &+ F_2(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_3\cdots\hat{a}_{2n+1}] \ ,\end{aligned} \tag{60}$$

$$\begin{aligned}\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_1\hat{a}_2\hat{a}_3\cdots\hat{a}_{2n+1}] &= F_3(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_3\cdots\hat{a}_{2n+1}] \\ &+ F_4(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_3\cdots\hat{a}_{2n+1}] \ ,\end{aligned} \tag{61}$$

$$\begin{aligned}\mathrm{Tr}[(1+\gamma^5)\hat{q}_+\hat{a}_1\hat{a}_2\hat{a}_3\cdots\hat{a}_{2n+1}] &= F_3(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_+\hat{a}_3\cdots\hat{a}_{2n+1}] \\ &+ F_4(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_3\cdots\hat{a}_{2n+1}] \ ,\end{aligned} \tag{62}$$

$$\begin{aligned}\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_1\hat{a}_2\hat{a}_3\cdots\hat{a}_{2n+1}] &= F_1(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_3\cdots\hat{a}_{2n+1}] \\ &+ F_2(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_+\hat{a}_3\cdots\hat{a}_{2n+1}] \ ,\end{aligned} \tag{63}$$

and so forth.

As an example, let us calculate

$$\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6] \ .$$

We take decomposition (21) written in the form

$$\begin{aligned}(1+\gamma^5)\hat{a}_6 &= (1+\gamma^5)\hat{q}_-(a_6q_+) + (1+\gamma^5)\hat{q}_+(a_6q_-)\\ &- (1+\gamma^5)\hat{e}_-(a_6e_+) - (1+\gamma^5)\hat{e}_+(a_6e_-) \ . \end{aligned} \tag{64}$$

Using (60) and (61), we get

$$\begin{aligned}&\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5]\\ =\ & F_1(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_3\hat{a}_4\hat{a}_5] + F_2(a_1,a_2)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_3\hat{a}_4\hat{a}_5]\\ =\ & F_1(a_1,a_2)\left\{F_1(a_3,a_4)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_5] + F_2(a_3,a_4)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_5]\right\}\\ +\ & F_2(a_1,a_2)\left\{F_3(a_3,a_4)\,\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_5] + F_4(a_3,a_4)\,\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_5]\right\}\\ =\ & [F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4)]\cdot 4(a_5q_-)\\ +\ & [F_1(a_1,a_2)F_2(a_3,a_4) + F_2(a_1,a_2)F_3(a_3,a_4)]\cdot 4(a_5e_-)\end{aligned}$$

i.e.

$$\begin{aligned}&\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5](a_6q_+)\\ =\ & 4[F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4)]\cdot(a_5q_-)(a_6q_+)\\ +\ & 4[F_1(a_1,a_2)F_2(a_3,a_4) + F_2(a_1,a_2)F_3(a_3,a_4)]\cdot(a_5e_-)(a_6q_+) \ .\end{aligned} \tag{65}$$

Making similar calculations for the remaining three items in decomposition (64), we obtain

$$\begin{aligned}&\mathrm{Tr}[(1+\gamma^5)\hat{q}_+\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5](a_6q_-)\\ =\ & 4[F_3(a_1,a_2)F_3(a_3,a_4) + F_4(a_1,a_2)F_2(a_3,a_4)]\cdot(a_5q_+)(a_6q_-)\\ +\ & 4[F_3(a_1,a_2)F_4(a_3,a_4) + F_4(a_1,a_2)F_1(a_3,a_4)]\cdot(a_5e_+)(a_6q_-) \ ,\end{aligned} \tag{66}$$

$$\begin{aligned}-\ &\mathrm{Tr}[(1+\gamma^5)\hat{e}_-\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5](a_6e_+)\\ =\ & -4[F_3(a_1,a_2)F_3(a_3,a_4) + F_4(a_1,a_2)F_2(a_3,a_4)]\cdot(a_5e_-)(a_6e_+)\\ -\ & 4[F_3(a_1,a_2)F_4(a_3,a_4) + F_4(a_1,a_2)F_1(a_3,a_4)]\cdot(a_5q_-)(a_6e_+) \ ,\end{aligned} \tag{67}$$

$$\begin{aligned}-\ &\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5](a_6e_-)\\ =\ & -4[F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4)]\cdot(a_5e_+)(a_6e_-)\\ -\ & 4[F_1(a_1,a_2)F_2(a_3,a_4) + F_2(a_1,a_2)F_3(a_3,a_4)]\cdot(a_5q_+)(a_6e_-) \ ,\end{aligned} \tag{68}$$

and finally

$$\begin{aligned}
&\frac{1}{2}\,\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6] \\
= \;& [F_1(a_1,a_2)F_1(a_3,a_4)+F_2(a_1,a_2)F_4(a_3,a_4)]F_1(a_5,a_6) \\
+ \;& [F_1(a_1,a_2)F_2(a_3,a_4)+F_2(a_1,a_2)F_3(a_3,a_4)]F_4(a_5,a_6) \\
+ \;& [F_3(a_1,a_2)F_3(a_3,a_4)+F_4(a_1,a_2)F_2(a_3,a_4)]F_3(a_5,a_6) \\
+ \;& [F_3(a_1,a_2)F_4(a_3,a_4)+F_4(a_1,a_2)F_1(a_3,a_4)]F_2(a_5,a_6)\ , \qquad (69)
\end{aligned}$$

that is we derive an expression consisting of eight terms only.

An expression for

$$\mathrm{Tr}[(1+\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6]$$

may be obtained from (69) by replacing all the functions with the complex-conjugate ones.

Having calculated

$$\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\cdots\hat{a}_{2n-1}\hat{a}_{2n}]\ ,$$

we may obtain an expression for

$$\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\cdots\hat{a}_{2n-1}\hat{a}_{2n}\hat{a}_{2n+1}\hat{a}_{2n+2}]$$

from the previous one by the following replacement:

$$F_1(a_{2n-1},a_{2n}) \to F_1(a_{2n-1},a_{2n})F_1(a_{2n+1},a_{2n+2})+F_2(a_{2n-1},a_{2n})F_4(a_{2n+1},a_{2n+2})\ , \qquad (70)$$

$$F_3(a_{2n-1},a_{2n}) \to F_3(a_{2n-1},a_{2n})F_3(a_{2n+1},a_{2n+2})+F_4(a_{2n-1},a_{2n})F_2(a_{2n+1},a_{2n+2})\ , \qquad (71)$$

$$F_2(a_{2n-1},a_{2n}) \to F_1(a_{2n-1},a_{2n})F_2(a_{2n+1},a_{2n+2})+F_2(a_{2n-1},a_{2n})F_3(a_{2n+1},a_{2n+2})\ , \qquad (72)$$

$$F_4(a_{2n-1},a_{2n}) \to F_3(a_{2n-1},a_{2n})F_4(a_{2n+1},a_{2n+2})+F_4(a_{2n-1},a_{2n})F_1(a_{2n+1},a_{2n+2})\ . \qquad (73)$$

Actually,

$$\begin{aligned}
&F_1(a_{2n-1},a_{2n}) = 2[(a_{2n-1}q_-)(a_{2n}q_+)-(a_{2n-1}e_+)(a_{2n}e_-)] \\
= \;& \frac{1}{2}\Big\{\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_{2n-1}](a_{2n}q_+)-\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_{2n-1}](a_{2n}e_-)\Big\} \\
\to \;& \frac{1}{2}\Big\{\mathrm{Tr}[(1+\gamma^5)\hat{q}_-\hat{a}_{2n-1}\hat{a}_{2n}\hat{a}_{2n+1}](a_{2n+2}q_+)-\mathrm{Tr}[(1+\gamma^5)\hat{e}_+\hat{a}_{2n-1}\hat{a}_{2n}\hat{a}_{2n+1}](a_{2n+2}e_-)\Big\} \\
= \;& 2\Big\{[F_1(a_{2n-1},a_{2n})(a_{2n+1}q_-)+F_2(a_{2n-1},a_{2n})(a_{2n+1}e_-)](a_{2n+2}q_+) \\
- \;& [F_1(a_{2n-1},a_{2n})(a_{2n+1}e_+)+F_2(a_{2n-1},a_{2n})(a_{2n+1}q_+)](a_{2n+2}e_-)\Big\} \\
= \;& F_1(a_{2n-1},a_{2n})F_1(a_{2n+1},a_{2n+2})+F_2(a_{2n-1},a_{2n})F_4(a_{2n+1},a_{2n+2})\ ,
\end{aligned}$$

and the procedure for the remaining three functions is quite similar.

Note that

$$\frac{1}{2}\,\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2] = 2(a_1a_2) = F_1(a_1,a_2) + F_3(a_1,a_2)\ . \tag{74}$$

Thus, formulae (74), (70) – (73) provide a very simple method of trace calculation (see **Appendix A**).

In numerical calculations, after fixing a reference system and choosing an orthonormal basis, for example, of the form

$$l_0^\mu = (1,\ 0,0,0)\ ,\ \ l_1^\mu = (0,\ 1,0,0)\ ,\ \ l_2^\mu = (0,\ 0,1,0)\ ,\ \ l_3^\mu = (0,\ 0,0,1)\ ,$$

we get

$$q_\pm^\mu = \frac{1}{\sqrt{2}}(1,\ \pm 1,0,0)\ , \qquad (aq_\pm) = \frac{1}{\sqrt{2}}(a_0 \mp a_x)\ ,$$

$$e_\pm^\mu = \frac{1}{\sqrt{2}}(0,\ 0,1,\pm i)\ , \qquad (ae_\pm) = -\frac{1}{\sqrt{2}}(a_y \pm i a_z)\ ,$$

and

$$F_1(a,b) = (ab) - (a_0b_x - a_xb_0) + i(a_yb_z - a_zb_y)\ , \tag{75}$$

$$F_3(a,b) = (ab) + (a_0b_x - a_xb_0) - i(a_yb_z - a_zb_y)\ , \tag{76}$$

$$F_2(a,b) = (a_0b_y - a_yb_0) + i(a_xb_z - a_zb_x) + (a_xb_y - a_yb_x) + i(a_0b_z - a_zb_0)\ , \tag{77}$$

$$F_4(a,b) = (a_0b_y - a_yb_0) + i(a_xb_z - a_zb_x) - (a_xb_y - a_yb_x) - i(a_0b_z - a_zb_0)\ , \tag{78}$$

that is we derive a number of the expressions well-suited for the implementation of a simple yet efficient computer algorithm.

Comparison of different methods for trace calculation is given in Tab. 1.

Table 1: Number of terms in the expression for traces

Number of γ-matrices, $2n$		2	4	6	8	10	12
Number of terms for algorithm (1)	$\mathrm{Tr}(\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n})$	1	3	15	105	945	10395
	$\mathrm{Tr}(\gamma^5\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n})$	0	1	15	210	3150	51975
Number of terms for algorithm (2)	$\mathrm{Tr}_4(\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n})$	1	3	15	105	693	4383
	$\mathrm{Tr}_4(\gamma^5\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n})$	0	1	6	33	180	1029
Number of terms for new method (70)–(74)	$\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n}]$	2	4	8	16	32	64

In case that traces of the form

$$\mathrm{Tr}(\gamma_\rho\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n+1})\,\mathrm{Tr}(\gamma^\rho\hat{b}_1\hat{b}_2\cdots\hat{b}_{2m+1})\ ,$$

need to be calculated, one may use the consequence of formula (20)

$$\gamma_\rho \times \gamma^\rho = \hat{q}_- \times \hat{q}_+ + \hat{q}_+ \times \hat{q}_- - \hat{e}_- \times \hat{e}_+ - \hat{e}_+ \times \hat{e}_- \ ,$$

or sum over index ρ using the classical methods, e.g.:

$$\begin{aligned} &\mathrm{Tr}(\gamma_\rho \hat{a}_1 \hat{a}_2 \cdots \hat{a}_{2n+1})\, \mathrm{Tr}(\gamma^\rho \hat{b}_1 \hat{b}_2 \cdots \hat{b}_{2m+1}) \\ &= 2\, \mathrm{Tr}[(\hat{a}_1 \hat{a}_2 \cdots \hat{a}_{2n+1} + \hat{a}_{2n+1} \cdots \hat{a}_2 \hat{a}_1) \hat{b}_1 \hat{b}_2 \cdots \hat{b}_{2m+1}] \ . \end{aligned}$$

Finally, we would like to note that for traces of four, six and eight Dirac γ-matrices we have directly compared the expressions for the traces obtained both by the new and classical methods:

- expressions for the new method were expanded by means of formulae (75) – (78) for functions F;
- in expressions for traces obtained by the classical method the scalar products and contractions of the Levi-Civita tensors with 4-vectors were written in terms of vector components.

The results obtained in different ways were identical.

Note that some other examples of applications of isotropic tetrads to the calculations in high energy physics may be found in [7].

A Expressions for traces

$$\frac{1}{2}\,\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2] = F_1(a_1,a_2) + F_3(a_1,a_2) \ ; \tag{79}$$

$$\begin{aligned} \frac{1}{2}\,\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4] &= F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4) \\ &+ F_3(a_1,a_2)F_3(a_3,a_4) + F_4(a_1,a_2)F_2(a_3,a_4) \ ; \end{aligned} \tag{80}$$

$$\begin{aligned} &\tfrac{1}{2}\ \mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6] \\ =\ & F_1(a_1,a_2)[F_1(a_3,a_4)F_1(a_5,a_6) + F_2(a_3,a_4)F_4(a_5,a_6)] \\ +\ & F_2(a_1,a_2)[F_3(a_3,a_4)F_4(a_5,a_6) + F_4(a_3,a_4)F_1(a_5,a_6)] \\ +\ & F_3(a_1,a_2)[F_3(a_3,a_4)F_3(a_5,a_6) + F_4(a_3,a_4)F_2(a_5,a_6)] \\ +\ & F_4(a_1,a_2)[F_1(a_3,a_4)F_2(a_5,a_6) + F_2(a_3,a_4)F_3(a_5,a_6)] \\ =\ & [F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4)]F_1(a_5,a_6) \\ +\ & [F_3(a_1,a_2)F_4(a_3,a_4) + F_4(a_1,a_2)F_1(a_3,a_4)]F_2(a_5,a_6) \\ +\ & [F_3(a_1,a_2)F_3(a_3,a_4) + F_4(a_1,a_2)F_2(a_3,a_4)]F_3(a_5,a_6) \\ +\ & [F_1(a_1,a_2)F_2(a_3,a_4) + F_2(a_1,a_2)F_3(a_3,a_4)]F_4(a_5,a_6) \ ; \end{aligned} \tag{81}$$

$$\begin{aligned}
\frac{1}{2} & \quad \mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\hat{a}_3\hat{a}_4\hat{a}_5\hat{a}_6\hat{a}_7\hat{a}_8] \\
= & \ [\ F_1(a_1,a_2)F_1(a_3,a_4) + F_2(a_1,a_2)F_4(a_3,a_4)] \\
\cdot & \ [\ F_1(a_5,a_6)F_1(a_7,a_8) + F_2(a_5,a_6)F_4(a_7,a_8)] \\
+ & \ [\ F_3(a_1,a_2)F_4(a_3,a_4) + F_4(a_1,a_2)F_1(a_3,a_4)] \\
\cdot & \ [\ F_1(a_5,a_6)F_2(a_7,a_8) + F_2(a_5,a_6)F_3(a_7,a_8)] \\
+ & \ [\ F_3(a_1,a_2)F_3(a_3,a_4) + F_4(a_1,a_2)F_2(a_3,a_4)] \\
\cdot & \ [\ F_3(a_5,a_6)F_3(a_7,a_8) + F_4(a_5,a_6)F_2(a_7,a_8)] \\
+ & \ [\ F_1(a_1,a_2)F_2(a_3,a_4) + F_2(a_1,a_2)F_3(a_3,a_4)] \\
\cdot & \ [\ F_3(a_5,a_6)F_4(a_7,a_8) + F_4(a_5,a_6)F_1(a_7,a_8)]\ ; \qquad (82)
\end{aligned}$$

and so forth.

$$\mathrm{Tr}[(1+\gamma^5)\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n}] = (\mathrm{Tr}[(1-\gamma^5)\hat{a}_1\hat{a}_2\cdots\hat{a}_{2n}])^*\ . \qquad (83)$$

B Functions $F_k(a,b)$

$$\begin{aligned}
F_1(a,b) &= 2[(aq_-)(bq_+) - (ae_+)(be_-)] = (ab) + \ G\begin{pmatrix} a & b \\ l_0 & l_1 \end{pmatrix} \ + i\ G\begin{pmatrix} a & b \\ l_2 & l_3 \end{pmatrix} \\
&= \frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{q}_+\hat{q}_-\hat{a}\hat{b}] = -\frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{e}_-\hat{e}_+\hat{a}\hat{b}]\ , \qquad (84)
\end{aligned}$$

$$\begin{aligned}
F_3(a,b) &= 2[(aq_+)(bq_-) - (ae_-)(be_+)] = (ab) - \ G\begin{pmatrix} a & b \\ l_0 & l_1 \end{pmatrix} \ - i\ G\begin{pmatrix} a & b \\ l_2 & l_3 \end{pmatrix} \\
&= \frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{q}_-\hat{q}_+\hat{a}\hat{b}] = -\frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{e}_+\hat{e}_-\hat{a}\hat{b}]\ , \qquad (85)
\end{aligned}$$

$$\begin{aligned}
F_2(a,b) &= 2[(ae_+)(bq_-) - (aq_-)(be_+)] = 2\ G\begin{pmatrix} a & b \\ e_+ & q_- \end{pmatrix} \\
&= -\ G\begin{pmatrix} a & b \\ l_0 & l_2 \end{pmatrix} \ + i\ G\begin{pmatrix} a & b \\ l_1 & l_3 \end{pmatrix} \ + \ G\begin{pmatrix} a & b \\ l_1 & l_2 \end{pmatrix} \ - i\ G\begin{pmatrix} a & b \\ l_0 & l_3 \end{pmatrix} \\
&= \frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{q}_-\hat{e}_+\hat{a}\hat{b}]\ , \qquad (86)
\end{aligned}$$

$$\begin{aligned}
F_4(a,b) &= 2[(ae_-)(bq_+) - (aq_+)(be_-)] = 2\ G\begin{pmatrix} a & b \\ e_- & q_+ \end{pmatrix} \\
&= -\ G\begin{pmatrix} a & b \\ l_0 & l_2 \end{pmatrix} \ + i\ G\begin{pmatrix} a & b \\ l_1 & l_3 \end{pmatrix} \ - \ G\begin{pmatrix} a & b \\ l_1 & l_2 \end{pmatrix} \ + i\ G\begin{pmatrix} a & b \\ l_0 & l_3 \end{pmatrix} \\
&= \frac{1}{4}\,\mathrm{Tr}[(1-\gamma^5)\hat{q}_+\hat{e}_-\hat{a}\hat{b}]\ . \qquad (87)
\end{aligned}$$

$$F_5(a,b) = F_1^*(a,b),\ F_6(a,b) = F_2^*(a,b),\ F_7(a,b) = F_3^*(a,b),\ F_8(a,b) = F_4^*(a,b). \quad (88)$$

References

[1] A.C. Hearn, *REDUCE user's manual. Version 3.5* (RAND publication, Santa Monica, 1993).

[2] J.S.R. Chisholm, Nuovo Cim. 30 (1963) 426.

[3] J. Kahane, J. Math. Phys. 9 (1968) 1732.

[4] H. Strubbe, Comput. Phys. Commun. 8 (1974) 1.

[5] J.A.M. Vermaseren, *Symbolic Manipulation with FORM* (CAN, Amsterdam, 1991).

[6] V.I. Borodulin, R.N. Rogalyov, S.R. Slabospitsky, IHEP-95-90 (Protvino, 1995), hep-ph/9507456.

[7] V.V. Andreev, Phys. Rev. D62 (2000) 014029, hep-ph/0101140.